

SAGITTARIUS B2 MAIN: A CLUSTER OF ULTRA-COMPACT HII REGIONS & MASSIVE PROTOSTELLAR CORES

JUN-HUI ZHAO¹ AND M. C. H. WRIGHT²

Draft version August 4, 2011

ABSTRACT

The ionized core in the Sgr B2 Main star-forming region was imaged using the Submillimeter Array archival data observed for the H26 α line and continuum emission at 0.86 millimeter with an angular resolution 0.3". Eight hyper-compact H26 α emission sources were detected with a typical size in the range of 1.6–20 $\times 10^2$ AU and electron density of 0.3–3 $\times 10^7$ cm⁻³, corresponding to the emission measure 0.4–8.4 $\times 10^{10}$ cm⁻⁶ pc. The H26 α line fluxes from the eight hyper-compact HII sources imply that the ionization for each of the sources must be powered by a Lyman continuum flux from an O star or a cluster of B stars. The most luminous H26 α source among the eight detected requires an O6 star that appears to be embedded in the ultra-compact HII region F3. In addition, ~ 23 compact continuum emission sources were also detected within the central 5" $\times$ 3" (~ 0.2 pc) region. In the assumption of a power-law distribution for the dust temperature, with the observed brightness temperature of the dust emission we determined the physical properties of the submillimeter emission sources showing that the molecular densities are in the range of 1–10 $\times 10^8$ cm⁻³, surface densities between 13 to 150 g cm⁻², and total gas masses in the range from 5 to $\gtrsim 200 M_\odot$ which are 1 or 2 orders of magnitude greater than the corresponding values of the Bonnor-Ebert mass. With a mean free-fall time scale of 2 $\times 10^3$ y, each of the massive protostellar cores are undergoing gravitational collapse to form new massive stars in the Sgr B2 Main core.

Subject headings: Galaxy: center — ISM: individual (Sgr B2) — HII region: radio lines — radio continuum: ISM — stars: formation

1. INTRODUCTION

The formation of massive stars is one of the challenging problems in astrophysics. Unlike their low-mass counterparts, massive stars are rare, and form in relatively deeply embedded massive molecular clouds on a much shorter time-scale (McLaughlin & Pudritz 1997; Osorio, Lizano & D'Alessio 1999; McKee & Tan 2002). High-mass stars are often formed in clusters (McKee & Ostriker 2007). After initial gravitational collapse of the natal clouds, multiple protostellar cores are formed through fragmentation of the gas. One of the fundamental theoretical questions is how the massive stars form in clusters. According to the competitive theory (Bonnell, Vine, & Bate 2004; Bonnell & Bate 2006; Bonnell 2008), mass clumps created from natal cloud gravitational collapse contain protostellar cores with small masses. These cores grow by accreting matter, competing with other cores, and form stars with many times their original mass. On the other hand, the direct gravitational collapse theory (Krumholz, McKee & Klein 2005) suggests that the protostellar cores created from fragmentation of the initial cloud gravitational collapse have sufficient mass to form individual high-mass to low-mass star systems in the subsequent collapse. Accretion from the parental cloud continues but does not substantially change their mass.

Strong radiation pressure from a newly formed massive star might halt infall, limiting the mass of stars

that can form (Kahn 1974; Wolfire & Cassinelli 1987; Larson & Starrfield 1971). Recent theoretical investigations suggest that the radiation pressure limit might be overcome in actual cases with complex, non-spherical infall geometries or high ram pressures in rotating disks (McKee & Tan 2002; Krumholz et al. 2009; Nakano 1989), or massive stars may form from stellar merging in a dense cluster (Bonnell, Bate, & Zinnecker 1998; Bonnell & Bate 2002). Numerical simulations suggest that gravitational instabilities cause the disk to fragment and form a massive companion to the primary (Krumholz et al. 2009), and consequently form binaries in a dense stellar system (Bonnell & Bate 2002). Furthermore, radiation feedback from massive stars in a cluster affects its subsequent fragmentation and consequently plays an important role in determining the stellar initial mass function (IMF) in a cluster (Krumholz et al. 2010).

In the past decade, good progress has been made in understanding massive young stellar clusters in the Galaxy. A recent review (Figer 2008) shows ten known Galactic clusters with masses $\leq 10^4 M_\odot$ with ages of a few million years. The Arches cluster in the Galactic center is the densest young cluster in the Galaxy and contains a large collection of massive stars (Figer 2005). Three out of ten (Quintuplet, Arches and Center) are located within the central 50 pc of the Galaxy, suggesting that the Galactic central region appears to prefer forming massive stars.

Located at a distance of 7.8 kpc (Reid et al. 2009) towards the Galactic center, Sgr B2, a giant molecular cloud (GMC) with a mass of 6 $\times 10^6 M_\odot$ (Goldsmith & Lis 1990) is one of the most active star-forming region in the Galaxy, radiating a to-

¹ Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA; jzhao@cfa.harvard.edu

² Department of Astronomy, University of California, Berkeley, Berkeley, CA 94720, USA

tal luminosity of $1 \times 10^7 L_{\odot}$ (Goldsmith & Lis 1990; Goldsmith et al. 1992). As the most luminous core among the several cores in this GMC, Sgr B2 Main is associated with numerous ultra-compact (UC) HII regions, suggesting the presence of a tight cluster of OB stars (Gaume & Claussen 1990; Gaume et al. 1995; De Pree et al. 1998). The observed molecular outflows and infalls suggest that on-going star formation activities are taking place (Qin et al. 2008; Rolfs, et al. 2010). Sgr B2 Main appears to be in a very young phase of forming a massive stellar cluster from the dense molecular core.

High-resolution observations using the Submillimeter Array (SMA)³ at submillimeter wavelengths can explore the detailed structure of the massive star forming core, providing useful clues on how massive stars form in a cluster.

The rest of this paper is organized as follows. In Section 2, we discuss the reduction and imaging process of the SMA archival data of Sgr B2 Main observed in 2007. Section 3 shows the results from the high-resolution observations of the H26 α line and continuum at 0.86 mm. Section 4 presents a model for the H26 α sources. In Section 5, the properties of the bright, compact dust emission clumps are determined and discussed. In Section 6, we discuss the kinematics in terms of ionized outflows/expansions and rotating disks. The early phase of massive star formation, and origin of the massive protostellar clumps are also discussed. In Section 7, we summarize the conclusions.

2. OBSERVATIONS AND DATA REDUCTION

The interferometer data for the H26 α line at $\nu_0 = 353.623$ GHz were acquired from the SMA archive, observed 2007 June 18, with the “very extended” array configuration in the upper-side band (USB). The reduction for the line data was made in Miriad (Sault, Teuben, & Wright 1995) following the reduction instructions for SMA data⁴. The bandpass calibration was made by applying a normalized average of all the data from the QSOs included in the observing run (J1229+020, J1733-130, J1743-038, J1751+096 and J2015+371). The flux density scale was determined by comparing observations of Callisto with the SMA planetary model. An angular size and brightness temperature of $1.5''$ and 120 K at the observing epoch were assumed. The complex gains of the data were primarily calibrated using the nearby QSO J1733-130 to obtain an image of the SgrB2 Main core. This image was used as an initial model to further correct residual errors in the visibilities using self-calibration.

To separate line and continuum emission, we used the Miriad task UVLIN. The line visibility data set was constructed by subtracting the continuum which was determined from line-free channels. We made images of the H26 α line with a channel width 3 km s^{-1} using a robustness weighting of 2 corresponding to natural weighting. The FWHM beam is $0.40'' \times 0.28''$ (PA=15°). The typical rms noise in a channel image is 0.1 Jy beam^{-1} .

A line-free continuum data set was also produced from the UVLIN program. We used both USB and LSB data to image the continuum emission with robustness weighting parameters 2 and -2 which correspond to natural and uniform weighting respectively. The FWHM beam with uniform weighting is $0.36'' \times 0.22''$ (PA=13°). The typical rms noise in the continuum images are 7.5 and 8.5 mJy beam⁻¹ for natural and uniform weighting, respectively. The shortest sampled spatial frequency in the visibility data is 30 k λ , corresponding to an angular size $\sim 6''$ – $7''$ for the H26 α line and continuum emission structure. The compact structure ($< 3''$) in the H26 α line and continuum emissions from the Sgr B2 Main core has been adequately sampled in the SMA observations.

We noticed a systematic offset of $0.3''$ in the phase center of the SMA images by comparing the positions of the UC HII regions G and F10.37 determined at 22.4 GHz with the VLA (Gaume et al. 1995). The offset could reflect residual errors in antenna positions and complex gain calibration due to the large distance between the calibration QSO J1733-130 and SgrB2 Main. We therefore imposed a shift of $0.3''$ for the phase center of the SMA data to align with the VLA image. The positional error of a compact source with $S/N = 6$ in the final SMA images with respect to the VLA coordinate frame is $\sim 0.03''$ (see below).

3. RESULTS

3.1. H26 α line emission

3.1.1. Distribution

Figure 1a shows the integrated H26 α line image of Sgr B2 Main made from line emission in the LSR velocity range between -100 and $+225 \text{ km s}^{-1}$. Eight sources with significant H26 α line emission ($> 6\sigma$ in the channel image) are labelled as H26 α -n, where n is numbered from 1 to 8. The source H26 α -1 has a small angular offset from the brightest continuum source (F3) at 22.4 GHz in the core of Sgr B2 Main while H26 α -8 has a larger angular distance from F3 (Gaume et al. 1995). The two relatively isolated bright line sources, H26 α -8 and H26 α -5, were used to align the 0.86 mm coordinate frame of our SMA images with the coordinate frame determined with the VLA at 22.4 GHz. The image shows the submillimeter positions of H26 α -8 and H26 α -5 agree with the 22.4 GHz positions of G and F10.37 within $\sim 0.01''$. H26 α -6 appears to coincide with F10.303. However, H26 α -1, the brightest line source, is located $0.17''$ northwest of F3. H26 α -6 coincides with F10.303. Two ultra-compact H26 α line sources (3 and 4) appear to be associated with F1. The hyper-compact H26 α sources 3 and 4 are located $\sim 0.15''$ northeast and southwest of F1, respectively. Also, H26 α -2 is located $0.2''$ southwest of F4. No significant H26 α emission associated with F2 has been detected. The positional offsets between the inner four H26 α emission sources (1,2,3 and 4) and their corresponding 22.4 GHz counterparts (F3, F4 and F1) appear to be significant, reflecting the fact that the free-free continuum emission at 22.4 GHz traces relatively lower-density ionized gas in the outer layer of the expanding HII gas or the ionized outflows, while the H26 α line traces the high-density ionized regions, possibly associated with the ionized disks or ultra- or hyper-compact ionized cores where the free-free emission appears to be optically thick

³ The Submillimeter Array is a joint project between the Smithsonian Astrophysical Observatory and the Academia Sinica Institute of Astronomy and Astrophysics and is funded by the Smithsonian Institution and the Academia Sinica.

⁴ <http://www.cfa.harvard.edu/sma/miriad>




FIG. 1.— **a**: SMA image of H26 α line emission at the rest frequency of 353.623 GHz, integrated from the velocity range of -100 to 125 km s $^{-1}$ (red contours and color) overlaid on the continuum image at 0.86 mm (yellow contours). The red contours are $F_P(\text{H26}\alpha) \times (4, 5, 7, 11, 17, 25, 35, 47, 61, 77$ and $95\%)$, and the yellow contours are $S_P(0.86\text{mm}) \times (-2, 2, 3, 5, 10, 15, 25, 35, 45, 55, 65, 75, 85,$ and $95\%)$, where the peak line flux $F_P(\text{H26}\alpha)$ and peak continuum flux density $S_P(0.86\text{mm})$ are 193 Jy beam $^{-1}$ km s $^{-1}$ and 2.88 Jy beam $^{-1}$, respectively. The FWHM beam is $0.40'' \times 0.28''$ (15°) denoted at bottom-right. The color wedge scales the H26 α line intensity in the units of Jy beam $^{-1}$ km s $^{-1}$ of the UC HII regions. The spectra from the eight brightest H26 α sources are also shown. The units of the vertical (line flux density) and horizontal (LSR velocity) axes in the spectral plots are Jy and km s $^{-1}$, respectively. The red cross in each spectral profile shows the peak line intensity (vertical bar) and the FWHM velocity width of a Gaussian profile determined from least-square fitting.

at 22.4 GHz.

3.1.2. Line broadening

The spectral profiles of the H26 α line from the eight brightest UC HII regions in the core of Sgr B2 Main are shown in Figure 1. H26 α -1, associated with F3, has a peak velocity $V_{\text{LSR}} = 64 \pm 1$ km s $^{-1}$ with a line width $\Delta V_{\text{FWHM}} = 40 \pm 1$ km s $^{-1}$ compared with $V_{\text{LSR}} = 68 \pm 2$ km s $^{-1}$ and $\Delta V_{\text{FWHM}} = 63 \pm 5$ km s $^{-1}$ derived from the VLA measurements of the H66 α line (De Pree et al. 1996). The peak velocity between the H26 α and H66 α line profiles shows a 2σ difference. The line width of the H66 α is much broader than that of H26 α . Using canonical values of $T_e = 1 \times 10^4$ K and $n_e = 1.1 \times 10^6$ cm $^{-3}$ for the typical UC HII regions in Sgr B2 Main (De Pree et al. 1998), for the VLA observations of the continuum emission at 7 mm, the thermal broadening ΔV_{th} contributes 21 km s $^{-1}$ to the line widths. The pressure broadening depends upon the principal quantum number N and electron density n_e ($\Delta V_P \propto n_e N^{7.4}$) (Brocklehurst & Leeman 1971;

Brocklehurst & Seaton 1972). We find that the pressure broadening ($\Delta V_P \sim 0.02$ km s $^{-1}$) is negligible for the H26 α line. Thus, the non-thermal broadening ΔV_{nth} including the Doppler motions of expansion, infall, outflows, rotation, shocks, and turbulence, obtained by subtracting the thermal broadening from the observed H26 α line width, is less than 34 km s $^{-1}$.

For the H66 α line, the pressure broadening for gas with $n_e = 1.1 \times 10^6$ cm $^{-3}$ accounts for 23 km s $^{-1}$. Subtracting ΔV_{th} , ΔV_{nth} and ΔV_P from the observed ΔV_{FWHM} for the H66 α line, $\sqrt{\Delta V_{\text{FWHM}}^2 - (\Delta V_{\text{th}}^2 + \Delta V_{\text{nth}}^2 + \Delta V_P^2)} \approx 43$ km s $^{-1}$, the residual line width, appears still to dominate the observed H66 α line width, suggesting that the electron density estimated from the 7 mm continuum data (De Pree et al. 1998) might be underestimated due to the large optical depth of the free-free emission from the higher density gas. If the residual line width is all due to pressure broadening, the observed large H66 α line width implies that a hyper-compact ionized core with an electron density $n_e > 2.5 \times 10^6$ cm $^{-3}$ is likely embedded in F3.




FIG. 2.— The image of 0.86-mm continuum emission observed with the SMA at a resolution of $0.36'' \times 0.22''$ (PA= 13°). The contours are $(-0.02, 0.02, 0.05, 0.15, 0.25, 0.35, 0.45, 0.55, 0.65, 0.75, 0.85, 0.95) \times 2.34 \text{ Jy beam}^{-1}$. The color wedge shows the intensity range in units of Jy beam^{-1} . The positions (cross) and the apparent shapes (ellipses) of the continuum sources are derived from Gaussian fitting.

3.2. Continuum Emission

The contours in Figure 1 show the continuum emission with the same FWHM beam ($0.4'' \times 0.28''$) as the H26 α line. The continuum flux densities corresponding to the H26 α line emission regions are summarized in column 8 of Table 2. A large fraction of the continuum emission at 0.86 mm appears to arise from a region outside the H26 α line sources, revealing an overall extended filamentary structure northwest to southeast. Figure 2 shows the details of the continuum emission at the higher resolution ($0.36'' \times 0.22''$). In addition to the UC HII regions observed at millimeter and centimeter wavelengths at the positions denoted with the red dots, we identified 23 continuum sources at submillimeter wavelengths which are listed in the bottom section of Table 2 and are named as Smm- with a sequential number (column 1) based on their angular distance $\Delta\theta$ from F3 (column 7). The corresponding sequential numbers of the Smm- sources are also marked in Figure 2. Using Gaussian fitting, we determined their positional offsets from F3 in RA and DEC (columns 2 and 3), deconvolved sizes (column 4) and

peak brightness and total flux densities (columns 5 and 6). The brightness temperatures of the sources are given in column 8. We note that the two brightest continuum cores are Smm-1 and Smm-4 with brightness temperatures 340 and 270 K, respectively. Smm-1 is located close to but with a significant offset $0.13'' \pm 0.01''$ from H26 α -1 which is associated with F3. Smm-4 is located $0.06'' \pm 0.02''$ southeast of its H26 α counterpart H26 α -4 in the F1 complex.

4. MODEL FOR UC HII REGIONS

On the assumption of isothermal homogeneous HII gas, a model has been developed for the low-frequency radio recombination lines (RRL) including non-LTE effects, such as stimulated emission by the background radiation (Shaver 1975). This model has been applied successfully to the high-frequency radio recombination lines (H30 α) at 231.9 GHz from the high-density gas in the minispiral within the circum-nuclear disk of the Galactic center (Zhao et al. 2010). The effects of the radiation from both synchrotron and dust sources seem to be negligible for

the H30 α line under the conditions within the central few parsecs of the Galaxy. In the massive star forming cores such as Sgr B2 Main, a considerable fraction of the continuum flux density at submillimeter wavelengths is from dust radiation that needs to be included in the model. Using a homogeneous isothermal model with temperatures of T_e and T_d for both HII gas and dust, the flux densities of hydrogen recombination line (S_L) and continuum (S_C) emission at radio to submillimeter wavelengths can be formulated assuming that the total dust is evenly distributed in three regions (each with an optical depth τ_d) along the line-of-sight — one third of the dust uniformly mixed with the ionized gas in a given HII region, and other two thirds are located in front of and behind the HII region, respectively. Equations (1) and (2) give the full solutions for the line and continuum flux densities from the three regions with no assumptions of small optical depths. Spectral fits to the data are shown in Figure 3. The results are not significantly different from an optically-thin dust approximation. This is because, in the centimeter to submillimeter wavelength range discussed in this paper, the dust emission towards the UC HII regions is indeed optically thin and we are not able to discern between the two models. However, at mid-IR (Spitzer) wavelengths, the dust becomes opaque and the dust attenuation towards the HII regions becomes important.

$$S_L = \frac{2k\nu^2}{c^2} \Omega T_e \left[\left(\frac{\tau_L/\beta_N + \tau_C}{\tau_L + \tau'_C} \right) \times \right. \\ \left. \left(1 - e^{-(\tau_L + \tau'_C)} \right) - \left(1 - e^{-\tau'_C} \right) \frac{\tau_C}{\tau'_C} \right] e^{-\tau_d} + \\ B(T_d)\tau_d\Omega \left[\frac{1 - e^{-(\tau_L + \tau'_C)}}{\tau_L + \tau'_C} - \frac{1 - e^{-\tau'_C}}{\tau'_C} \right] e^{-\tau_d} + \\ B(T_d)(1 - e^{-\tau_d})\Omega \left[e^{-\tau_L} - 1 \right] e^{-(\tau'_C + \tau_d)}, \quad (1)$$

and

$$S_C = \frac{2k\nu^2}{c^2} \Omega T_e \left(1 - e^{-\tau'_C} \right) \frac{\tau_C}{\tau'_C} e^{-\tau_d} + \\ B(T_d)\tau_d\Omega \frac{1 - e^{-\tau'_C}}{\tau'_C} e^{-\tau_d} + \\ B(T_d)(1 - e^{-\tau_d})\Omega \left[1 + e^{-(\tau'_C + \tau_d)} \right], \quad (2)$$

where k is Boltzmann's constant, c is the speed of light, and the total continuum optical depth in the HII region $\tau'_C = \tau_C + \tau_d$ includes both the free-free and dust contributions. The recombination line (τ_L) and free-free continuum (τ_C) optical depths of the radiation from an HII region are given by Equations (A3) and (A4) in Appendix A, respectively. The values of the population departure coefficients (b_N and β_N) were calculated using the non-LTE code of Gordon & Soroichenko (2002) based on the analysis of hydrogen recombination lines

at wavelengths from radio to submillimeters (Walmsley 1990). The quantity $B(T_d)$ is the Planck function with a dust temperature T_d . In Equation (1), the first term associated with $\frac{2k\nu^2}{c^2} \Omega T_e$ is the line emission from the HII gas attenuated by the foreground dust with optical depth τ_d ; the second term associated with $B(T_d)\tau_d\Omega$ accounts for the absorption of the internal dust radiation by the HII gas if $\tau_L \geq 0$, or the line emission stimulated by the internal dust radiation if $\tau_L < 0$; and the third term associated with $B(T_d)(1 - e^{-\tau_d})\Omega$ is the absorption of background dust radiation by the HII gas if $\tau_L \geq 0$, or the line emission stimulated by the background dust radiation if $\tau_L < 0$. On the other hand, for an isolated dust source, the dust continuum flux density is

$$S_d = B(T_d)(1 - e^{-\tau_d})\Omega. \quad (3)$$

The optical depth τ_d of dust continuum radiation can be described as

$$\tau_d = \kappa_0 \left(\frac{\nu}{\nu_0} \right)^\beta M_d D^{-2} \Omega^{-1}, \quad (4)$$

where $\kappa_\nu = \kappa_0(\nu/\nu_0)^\beta$ is the dust opacity per unit dust mass, M_d is dust mass, D is the distance, and $B_\nu(T_d)$ is the Planck function with a dust temperature of T_d . In the calculations throughout this paper, we adopted $\kappa_0 = 1.06 \text{ cm}^2 \text{ g}^{-1}$ at the reference frequency $\nu_0 = 230 \text{ GHz}$ calculated by Ossenkopf & Henning (1994) for high-density gas $n_H = 10^7 \text{ cm}^{-3}$ using the standard MRN model (Mathis, Rumpl & Nordsieck 1977) with thin ice mantles. The physical parameters for a given HII region can be determined by fitting an isothermal, homogeneous model to the data at the submillimeter wavelengths of this paper and the data taken from previous observations at radio and millimeter wavelengths as discussed in the following section.

4.1. Radio continuum spectrum & RRL

F3 is the brightest UC HII region in the Sgr B2 Main core as observed at wavelengths of 2, 1.3 and 0.7 cm (Gaume & Claussen 1990; Gaume et al. 1995; De Pree et al. 1998). **F3**, which is associated with the brightest H26 α line source, H26 α -1, was detected at 2 cm with a peak flux density $0.138 \text{ Jy beam}^{-1}$ (with a circular beam of $0.3''$), and a total flux density 0.48 Jy in an extended region of size $0.7''$, FWHM (Gaume & Claussen 1990). With a size $0.5'' \times 0.5''$, FWHM, the peak intensity and total flux density at 1.3 cm are $0.22 \text{ Jy beam}^{-1}$ and 0.9 Jy (Gaume et al. 1995). Five HII components have been detected at high-angular resolution ($0.049'' \times 0.079''$) with the VLA at 7 mm (De Pree et al. 1998). The authors show that the major component F3-d has a shell- or ring-like morphology. Observations with the Hat-Creek interferometer array at 3.5 mm, Liu & Snyder (1999) showed that the peak emission of $1.51 \text{ Jy beam}^{-1}$ (FWHM = $1'' \times 0.5''$) is coincident with F3 corresponding to a total flux density of 4.38 Jy for F1, F2, F3 and F4. The high-resolution ($0.64'' \times 0.31''$) SMA image at 1.3 mm shows that the peak emission of 1.8 Jy coincides with F3. The H66 α line has been detected with $1.1'' \times 1.6''$ resolution at $V_{\text{LSR}} = 68 \pm 2 \text{ km s}^{-1}$ showing a line-to-continuum (L/C) ratio of $3 \pm 0.3 \%$ and $\Delta V_{\text{FWHM}} = 63 \pm 5 \text{ km s}^{-1}$ determined from integrated




FIG. 3.— Models for the UC HII regions F3, F4, F1A, F1B, F10.37, F10.303, F10.39 and G with the constraints from the observed radio continuum flux densities and the fluxes of hydrogen recombination lines at radio and submillimeter wavelengths. The red circles denote the integrated flux ($1.064 S_p * \Delta V_{\text{FWHM}}$) of the hydrogen recombination lines at H26 α (from the SMA measurements of this paper), H52 α and H66 α from the VLA measurements (De Pree et al. 1996). The black dots are the flux densities determined at 2, 1.3 cm and 7 mm with the VLA and 3 mm with the Hat Creek interferometer array and 1.3 and 0.86 mm with the SMA (see Section 4.1, the 1.3 mm data from Zhu 2010, personal communication). The red curves indicate the fluxes of the hydrogen recombination lines from the best fitted model (see section 4) to the observed data. The black curves indicate the continuum spectral energy distribution (SED) resultant from the free-free (flat dashed curves corresponding to the first term in Equation (2)) and dust (steep rising curves corresponding to the second term in Equation (2)) emissions calculated from the best fitted model.

line profile (De Pree et al. 1996). At the lower resolution $2.9'' \times 1.6''$, the H52 α line is towards the F cluster at $V_{\text{LSR}} = 57 \pm 2 \text{ km s}^{-1}$ with L/C of $14 \pm 2 \%$ and $\Delta V_{\text{FWHM}} = 59 \pm 2 \text{ km s}^{-1}$ (De Pree et al. 1996). De Pree et al. (1996) noticed that the LTE electron temperature (T_e^*) derived from the high L/C value of the H52 α line is significantly lower by a factor of 2 than that derived from the H66 α line under the assumption of optically thin and LTE gas. If the L/C ratios are uniform across the F complex, the fluxes of the H66 α and H52 α lines from the dominant source F3 can be estimated from the L/C values and the continuum flux densities at the corresponding wavelengths.

For the other seven regions associated with the H26 α line source, the continuum flux densities at 2, 1.3 and 0.7 cm used in this paper are from Gaume & Claussen (1990), Gaume et al. (1995) and De Pree et al. (1998). The H66 α line flux from the UC HII region G (H26 α -8) is estimated from the L/C ratio determined by De Pree et al. (1996) and the continuum flux density

of Gaume et al. (1995).

Figure 3 shows the observed spectra of the continuum (black dots) and hydrogen recombination lines (red dots) at radio and submillimeter wavelengths.

In order to fit to both the continuum and hydrogen recombination line data obtained from the high-resolution observations at wavelengths in the range between 2 cm to 0.86 mm, we considered a few possible models. A model with an LTE approach and a single isothermal, homogeneous HII component was rejected since the observed large H26 α line flux requires large free-free continuum optical depths ($\tau_C > 1$) at longer millimeter and centimeter wavelengths for a single component model. At these wavelengths, the dust absorption ($\tau_d \ll 1$) is negligible. Then, the predicted line fluxes at centimeter wavelengths for larger quantum-number transitions are much smaller than the observed values because of the exponential drop in the LTE line flux due to the large free-free optical depth $S_L^* = \frac{2k\nu^2}{c^2} \Omega T_e (1 - e^{-\tau_L}) e^{-\tau_C}$.

The observed line flux densities at centimeter wavelengths could be fitted with a single isothermal, homogeneous component by adjusting n_e and T_e if the stimulating effect in a non-LTE approach is considered. However, the flat continuum spectra at centimeter wavelengths seen in all the eight HII sources (Figure 3) place a critical restriction on a model with a single isothermal, homogeneous component, requiring an additional lower-density component. Thus, we fit each of the eight observed hyper-compact H26 α sources and the UC HII regions surrounding them with two components corresponding to high- and low-density ionized gas with small and large sizes, A and B, respectively. Further assumptions used in the two-component model are the volume filling factors $f_V = 0.1$ and 0.5 for components, A and B, respectively. The effect of shadowing between the two components is negligible.

4.2. Hyper-compact HII component

Figure 3 shows the results of model fitting the observed hydrogen recombination line (red curves) and continuum flux densities (black curves) from each of the eight UC HII regions with two isothermal, homogeneous HII components A and B. The best fitted parameters are summarized in Table 3. The first row gives the size in units of 10^3 AU. The following rows under the category of HII gas properties summarize the derived parameters including the temperature T_e , density n_e , volume filling factor f_V , the departure coefficients b_N and β_N , the optical depths of the H26 α line τ_L (H26 α), the free-free continuum τ_C (0.86 mm) at 0.86 mm, the fractional contribution $\zeta_{H26\alpha}$ to the observed H26 α line flux from each component, the H66 α line τ_L (H66 α), and the free-free continuum τ_C (13 mm) at 13 mm. Components A have small size (160 to 2000 AU), high electron density (3 to 33×10^6 cm $^{-3}$), relatively lower temperature (5 to 9×10^3 K). The A components are typically 10 times smaller but 100 times denser than typical UC HII regions, 10^{17} cm and 10^4 cm $^{-3}$, (Churchwell 2002), representing a class of hyper-compact HII components (Churchwell 2002) present in the Sgr B2 Main core. The hyper-compact HII components could arise from plausible ionized disks with emission measure (EM) in the range 0.4 to 8.4×10^{10} cm $^{-6}$ pc, accounting for the most of the H26 α line flux, *e.g.* $\sim 94\%$ for F3. The hyper-compact HII components in all the HII cores show a negative optical depth for the H26 α line, suggesting the presence of stimulated emission by free-free continuum emission from the hyper-compact HII region itself. For F3, the H26 α line is enhanced by a factor of 1.4 due to a weak stimulating process within the hyper-compact HII component (A) while the enhancement factors (S_L/S_L^*) are 1.7 and 4 for the H26 α lines from G and F10.39 which show larger negative line optical depths of -0.71 and -1.3 , respectively. At 13 mm, the continuum component A becomes optically thick while the H66 α line is still optically thin, τ_C (13mm) = 9.5 and τ_L (H66 α) = 0.13 for F3. Component A appears to make a little contribution to the total line flux of the H66 α transition ($\sim 20\%$ for F3).

4.3. Ultra-compact HII component

The continuum emission at longer wavelengths from each of the UC HII regions shows a shallow rising spec-

tra towards short wavelengths, indicating the presence of a larger HII component which might result from an ionized stellar wind or expansion of an ionized nebula. Component (B) with a relatively large size (1200 to 4000 AU), lower density (0.1 to 3.5×10^5 cm $^{-3}$), and higher temperature (10 to 17×10^3 K) is needed for the extended ionized gas with relatively smaller emission measures of $\leq 1 \times 10^9$ cm $^{-6}$ pc. $T_e = 17,000$ K for the UC HII component in F3 (H26 α -1) appears too high for the HII regions with metallicities in the Galactic center. The over-estimated T_e from our fitting might be caused by the under-estimate of the H66 α and H52 α line fluxes at 1.3 and 0.7 cm due to large line broadening and limited bandwidth coverage of the old VLA system. For a canonical value of $T_e = 10,000$ K, the corresponding H66 α and H52 α line fluxes require 50% more than the values used in this paper. The physical parameters of the B components fall into the category of UC HII region in Churchwell (2002). The continuum emission from the B components is optically thin at longer millimeter wavelengths, providing a flat spectrum ($\propto \nu^{-0.1}$) in contrast to the steep rising spectrum ($\propto \nu^2$) of component A. With its larger size, the free-free emission from the lower density gas (component B) dominates the continuum flux densities at 13 mm. Superposition of the spectra from the two components (A and B) results in a spectrum slowly rising at long wavelengths and turning over to flat at short millimeter and sub-millimeter wavelengths.

4.4. Dust component

The composite spectra (dashed flat curves in Figure 3) of components A and B appear to fit the observed continuum flux densities at millimeter and centimeter wavelengths. In the submillimeter, the observed flux densities show a significant excess emission from the UC HII region over the ones predicted from the isothermal, homogeneous ionized gas model with two-density components, suggesting that dust radiation becomes significant in the continuum flux density. In order to evaluate the contribution of dust radiation in the UC HII regions, the terms involving the dust radiation $B(T_d)$ in Equations (1) and (2) are added to the B component model with the dust distribution as assumed. The flux density (S_d) of the dust radiation is determined using Equations (3) and (4) with an assumed dust temperature in the HII region equal to the peak temperature in the Sgr B2 Main core, $T_d \sim 500$ K, (Lis & Goldsmith 1990) and a power-law index for the dust opacity $\beta = 1.5$ (Goldsmith & Lis 1990; Lis & Goldsmith 1990). The observed brightness temperatures of few times 10^2 K for the continuum emission from several central dense clumps also suggest a high dust temperature for the UC HII regions in Sgr B2 Main (also see the discussion in Section 5 for the high dust temperature).

We show that, except for G, the dust radiation in the other seven regions makes a contribution to the observed continuum flux density at 0.86 mm comparable or larger than that from free-free emission (see Figure 3). In general, the dust in each of the UC HII regions is optically thin, $\tau_d \sim 0.1$, with a total gas mass $\leq 11 M_\odot$. The dust optical depth and the gas mass (M_{tot}) needed in each of the UC HII regions are summarized in Table 3.

Finally, the models with two independent HII com-

ponents mixed with dust involved a set of ten free parameters, namely $2 \times (n_e, T_e, R, f_V, \text{ and } T_d)$. For F3, we have collected a total of nine measurements in line and continuum flux densities from high-resolution observations at centimeter-submillimeter wavelengths. Adding the two determined angular sizes for the components A and B, we have a total of eleven measurements to constrain the model reasonably well. However, for the rest of the sources, the models need to be confined with further high-resolution observations of both continuum and line emissions at wavelengths from centimeters to sub-millimeters.

4.5. Lyman continuum

The hyper-compact HII components detected with H26 α lines appear to require most of the flux of Lyman continuum photons from the ionizing stars to maintain their ionization. The H26 α line flux from hyper-compact sources would be excellent tracers for newly formed massive stars. Given an observed FWHM linewidth of ΔV_{FWHM} that is dominated by Doppler broadening ΔV_D in the H26 α line profile, the ionizing photon flux required for the observed H26 α line flux (S_L) can be estimated from Equation (A8) that was generally derived for a radio recombination line $N + 1 \rightarrow N$,

$$N_{\text{Lym}} = 7.8 \times 10^{46} \left(\frac{S_L \Delta V_D}{\text{Jy km s}^{-1}} \right) \left(\frac{\nu_{\text{H26}\alpha}}{\nu} \right) \left(\frac{D}{8 \text{ kpc}} \right)^2 \times \left(\frac{T_e}{10^4 \text{ K}} \right)^{1.5} \left(\frac{\alpha_B(T_e)}{\alpha_B(10^4 \text{ K})} \right) \Psi_{26}^{-1} \text{ ph s}^{-1} \quad (5)$$

where $\alpha_B(T_e)$ is the total recombination coefficient to excited levels (Hummer & Seaton 1963), and the function Ψ_{26} provides a correction for the effects of both non-LTE and optical-depth. The values of Ψ_{26} are given in Table 3 for each case. For the hyper-compact HII components (A) in Sgr B2 Main, Ψ_{26} is in the range ~ 1 (H26 α -6) to ~ 2 (H26 α -7). For the UC HII components (B) where the H26 α line is optically thin, $\Psi_{26} \approx b_{26}$, a correction for the factor due to the lower quantum-number levels that are underpopulated with respect to the LTE in the cases of lower electron density. The electrons at the level of $N = 26$ appear to be underpopulated by 20% to 30% with respect to the LTE for the UC HII regions in Sgr B2 Main.

In addition, for H26 α -1 (F3) assuming $T_e = 9,000 \text{ K}$, the value $N_{\text{Lym}} = 1.43 \times 10^{49} \text{ ph s}^{-1}$ is inferred from the observed H26 α line flux 280 Jy km s^{-1} using Equation (5) with $\Psi_{26} = 1.3$ while $N_{\text{Lym}} = 2.66 \times 10^{49} \text{ ph s}^{-1}$ is derived from the continuum flux density $S_C = 2.9 \text{ Jy}$ at 353.6 GHz using Equation (4) given by Wilcots (1994). The value determined from the observed continuum flux density surpasses the value determined from H26 α line flux by 46%. Our detailed model fitting gives the free-free flux density of 1.61 Jy suggesting the fraction of dust contribution to the continuum is 44% at 353.6 GHz, in good agreement with above assessment. Considering optically thin of the line emission and nearly no attenuation and contamination by the dust at 353.6 GHz (0.85 mm), it appears to be an excellent way to determine the N_{Lym} and the free-free flux density using the H26 α line flux. Therefore, comparing the observed continuum flux den-

sity with the free-free flux density determined from the H26 α line flux, one can give a good assessment of the fraction of dust contribution to the continuum.

4.6. Ionizing stars

Based on the derived properties for both A and B components, we estimated the fluxes of the Lyman continuum photons (N_{Lym}) from the newly formed ionizing stars. In each of the UC HII regions, the hyper-compact component (A) appears to require more flux in Lyman continuum photons than their larger but lower-density counterpart (component B) for the maintenance of the ionization (see Table 3). A total flux in Lyman continuum photons ($N_{\text{Lym}}^{\text{A+B}}$) is evaluated by the addition of the individual fluxes required for components A and B. Assuming that $N_{\text{Lym}}^{\text{A+B}}$ accounts for all the Lyman continuum photons from a single early-type, zero-age-main-sequence (ZAMS) star in each of the UC HII regions with no significant leakages, *i.e.* the UC HII regions are internally ionized (De Pree et al. 1998), the type of massive stars required for each of eight HII complexes with a hyper-compact H26 α component is inferred on the basis of the stellar atmosphere model computed by Panagia (1973) and listed in Table 3. We note that the Lyman continuum photon fluxes from Panagia (1973)'s stellar atmosphere model were underestimated by 26% to 66% for O6 to B0 type stars, respectively, as compared to those computed from Vacca, Garmany & Shull (1996)'s model. In good agreement with De Pree et al. (1998), we also find that at least an O6 star is required for the brightest H26 α source (H26 α -1) in F3 and an O8.5 star for H26 α -2 in F4.

The complex F1, including H26 α -3 (O7), H26 α -4 (O9.5) and H26 α -6 (O9), has been resolved by the VLA with resolution of $0.049'' \times 0.079''$ into at least seven compact components (De Pree et al. 1998). The authors suggest that the complex requires a group of seven early-B and late-O type stars (B0 to O8.5) to maintain the ionization. The flux of Lyman continuum photons (N_{Lym}) inferred from our analysis of the H26 α line for the hyper-compact sources in F3 and F1 appear to be considerably greater than the values inferred from their 7 mm-counterparts based on the VLA continuum observations (De Pree et al. 1998). The difference occurs because the hyper-compact components H26 α -1 and H26 α -3 in F3 and F1 are optically thick in free-free emission at 7 mm, $\tau_C \approx 2$, inferred from the analysis above on the basis of H26 α observations. The hyper-compact core appears to be deeply embedded in the optically thick region at 7 mm. From the VLA flux densities at 7 mm, the flux of Lyman continuum photons has possibly been underestimated due to missing the contribution from the hyper-compact ionized core. Thus, an O7 star might be needed to maintain the ionization of hyper-compact H26 α -3 in the HII complex of F1.

The above argument is also valid for the hyper-compact HII components H26 α -5 (O9), H26 α -7 (O9.5) and H26 α -8 (O8.5). These hyper-compact components appear to be embedded in the relatively isolated UC HII regions F10.37, F10.39 and G, respectively. In fact, the continuum optical depths inferred for the three hyper-compact HII components are greater than the values derived for the rest of hyper-compact HII components (see

Table 3).

As discussed above (see Figure 3), the H26 α line from the three components appears to be enhanced by stimulated emission from continuum emission in the hyper-compact HII gas surrounding the newly formed O stars. The actual flux of the ionizing photons is reduced by a factor of 4 in H26 α -7 as compared to an LTE source. Thus, the effect of stimulated line emission helps detection of H26 α line from a region ionized by an O9.5 among the eight H26 α line sources in SgrB 2 Main.

Given a hyper-compact HII region with $T_e = 1 \times 10^4$ K, $\Delta V_D = 30$ km s $^{-1}$ without line stimulation, the 3σ (0.3 Jy beam $^{-1}$) detection limit for the H26 α line imposed by the SMA data used in this paper gives a limit of $N_{\text{Ly}\alpha} = 7.1 \times 10^{47}$ ph s $^{-1}$, or $\log(N_{\text{Ly}\alpha}) = 47.85$, on the flux of ionizing photons, corresponding to an O9.5 ZAMS star. No significant detections of the H26 α line with the SMA towards F2 are consistent with four B0 stars inferred from the 7-mm continuum observations (De Pree et al. 1998).

4.7. Ages

The ages of the newly formed O stars in Sgr B2 Main can be assessed by the dynamic time-scale of the larger HII components (B) as the time of sound wave traveling from the initial ionization front at a radius close to the Strömgren radius (R_S) (Strömgren 1939) as suggested by Spitzer (1978) and Garay & Lizano (1999). Using the Equation (3) given in Shi, Zhao, & Han (2010b) and the isothermal sound speed $C_S = \sqrt{kT_e/m_H}$, we calculated the dynamical age of the HII components (B) and listed the values in Table 3, ranging between the oldest one, 1.3×10^3 y for F3 (H26 α -1) and the youngest one, 0.4×10^3 y for F10.37 (H26 α -7). If there is any external pressure produced by the swept-up ISM as the nebula expands, the sound crossing time will give an underestimate of the age. The O type stars in Sgr B2 Main appear to be clustered around a thousand years ago.

5. A CLUSTER OF PROTOSTELLAR CORES

The dust emission from Sgr B2 Main has been resolved into at least twenty three components, showing that a cluster of protostellar cores are present in this region in addition to the newly formed O and B type stars suggested by SMA observations of the H26 α line and VLA observations of the radio continuum emission and H52 α and H66 α lines. From our interferometer observations, the brightness temperature (T_b) of the continuum emission from individual components can be determined under the Rayley-Jeans approximation,

$$T_b(\nu) \approx 13.6 \lambda_{\text{mm}}^2 \left[\frac{S_d}{Jy} \right] \theta_{\text{FWHM}}^{-2} \text{ K}, \quad (6)$$

where S_d is the dust flux density at wavelength, λ from the region with the geometric mean of the angular size θ_{FWHM} in arcsec. Located near $(0.13'' \pm 0.01'' \text{ north})$ the brightest H26 α line source in F3, Smm-1 has the highest brightness temperature of 340 K. The contribution to the brightness temperatures in the continuum cores from the free-free continuum emission is less than 10%. Thus, the brightness temperature from dust emission is greater than 300 K in the center, suggesting that high dust temperature is present in Sgr B2 Main.

From Equations(3) and (5), for a given dust temperature T_d , we can determine the optical depth (τ_d) for each of the dust cores from $T_b(\nu)$:

$$\tau_d(\nu) = \log \left[\frac{T_d}{T_d - T_b(\nu)} \right] \quad (7)$$

The dust temperature (T_d) is affected by the increasing luminosity in the inner region (~ 0.1 pc) of the Sgr B2 Main core where at least eight newly formed O stars are suggested by the H26 α line sources. We determined the physical properties of the proto-stellar cores using a power-law distribution for the mean dust temperature T_d as function of the radial distance r from the center. The distribution of T_d is modeled by Scoville & Kwan (1976) and Wolfire & Cassinelli (1987) assuming that the total emissivity of the collection of grains at that temperature equals the total radiative energy that is absorbed by the grains (Scoville & Kwan 1976; Wolfire & Cassinelli 1986),

$$T_d = T_{\text{in}} \left(\frac{r}{r_{\text{in}}} \right)^{-\Gamma} \quad (8)$$

where T_{in} is the dust temperature at $r_{\text{in}} = 0.01$ pc, the radial distance between Smm-1 and H26 α -1 (O6 star) assuming a mean projection angle of 45°. Depending on the dust emissivity (Q_ν), the central stellar luminosity is insensitive to the exponent β of the dust emissivity power-law ($\Gamma = 2/5$ or $1/3$ for $\beta = 1$ or 2 , respectively).

We used a linearly interpolated value of $\Gamma = 0.37$ for $\beta = 1.5$ in the modeling. Assuming dust heated primarily by the central O6 star with a luminosity of $2.5 \times 10^5 L_\odot$, $T_{\text{in}}(r_{\text{in}} = 0.007 \text{ pc}) \approx 400$ K is inferred from the structure of the dust temperature derived by Scoville & Kwan (1976). On the other hand, a larger value of $T_{\text{in}}(r_{\text{in}} = 0.007 \text{ pc}) \approx 630$ K is calculated by Lis & Goldsmith (1990) in their detailed models assuming a total luminosity up to $2.3 \times 10^7 L_\odot$ from the stars distributed in the core in Sgr B2 Main and exponent of $\beta = 1.5$ in the dust emissivity power-law. Because of high infrared opacity in the inner region of the core, the radiation from the outer region does not affect the dust temperature at the center. The observed brightness temperature, $T_b = 300$ K from the dust in Smm-1 indicates the dust temperature $T_d \approx 470$ K if $\tau_d \approx 1$, which is consistent with the kinetic temperature of the absorbing gas towards Sgr B2 Main (Cernicharo, Goicoechea, & Pardo 2006; Qin et al. 2008). We note that the dust sublimation radius R_{sub} by heating from the central star can be determined by balancing the dust absorption of the UV and visible radiation from the star with the re-emission at IR from the dust grains at its sublimation temperature $T_{\text{sub}} = 1800$ K (Wolfire & Cassinelli 1986). For an O6 star, $R_{\text{sub}} = 2.9 \times 10^{15}$ cm, is about an order magnitude smaller than r_{in} and in the zone between the two radial distances, the dust substantially cools down due to infrared radiation. In the following calculation, we take $T_{\text{in}} = 500$ K in the power-law exponent of $\Gamma = 0.37$ for the dust temperature. Using Equation (8), we calculate T_d in the range 500 K for Smm-1 near the O6 star to 170 K for Smm-23 about 0.14 pc away. Using Equations (6), (7) and (8), we find that dust optical depth lies between 0.3 and 3.0 at 0.86 mm for these protostellar cores, with unity for the brightest core Smm-1. Table 4 summarizes

the optical depths (τ_d)

From Equations (9), (10) and (11), we can calculate the total gas masses (M), surface density (Σ) and molecular number density (n_H) in the individual protostellar cores assuming a mass ratio of gas-to-dust $g = 100$ and a mean mass per molecule, $\mu = 2.3$,

$$M = 12.0 \left[\frac{\nu}{230\text{GHz}} \right]^{-\beta} \log \left[\frac{T_d}{T_d - T_b(\nu)} \right] \times D_{\text{kpc}}^2 \theta_{\text{FWHM}}^2 M_{\odot}, \quad (9)$$

$$\Sigma = 94.3 \left[\frac{\nu}{230\text{GHz}} \right]^{-\beta} \log \left[\frac{T_d}{T_d - T_b(\nu)} \right] g \text{ cm}^{-2}, \quad (10)$$

and

$$n_H = 16.4 \times 10^8 \left[\frac{\nu}{230\text{GHz}} \right]^{-\beta} \log \left[\frac{T_d}{T_d - T_b(\nu)} \right] \times D_{\text{kpc}}^{-1} \theta_{\text{FWHM}}^{-1} \text{ cm}^{-3}. \quad (11)$$

Table 4 lists the derived values for M , Σ and n_H in columns (6), (7) and (8). The model predicts a mass $143 M_{\odot}$ for Smm-1, while the largest mass of $223 M_{\odot}$ is associated with Smm-4 near F1 (H26 α -3). The enclosed gas masses (M) of the individual dense cores range 5 to $223 M_{\odot}$. The molecular surface density is in the range 13 to $150 g \text{ cm}^{-2}$. The molecular number density is in the range $1 \times 10^8 - 1 \times 10^9 \text{ cm}^{-3}$, which appears to be consistent with the ambient density of $10^7 - 10^8 \text{ cm}^{-3}$ suggested by De Pree et al. (1998) if ionized gas in the UC HII regions and the ambient molecular gas are in pressure equilibrium (Xie et al. 1996).

In order to evaluate whether the individual dust cores are subject to inevitably undergoing gravitational collapse to form stars, we calculated the Bonnor-Ebert mass (M_{BE}), the maximum mass that a dense core remains in hydrostatic equilibrium (Bonnor 1956; Ebert 1957), and free fall time scale (t_{ff}),

$$M_{\text{BE}} = 1.18 C_S^3 G^{-3/2} \rho^{-1/2} \quad (12)$$

$$t_{\text{ff}} = \frac{1}{4} \sqrt{\frac{3\pi}{2G\rho}} \quad (13)$$

where C_S , G and ρ are the sound speed, gravitational constant and mass density of the core. The derived values for M_{BE} and t_{ff} are listed in columns 9 and 10 of Table 4. The masses of the protostellar cores in the central region of Sgr B2 Main are 1 to 2 orders of magnitude greater than their corresponding Bonnor-Ebert mass. Free-fall time scales of $1 - 3 \times 10^3 \text{ y}$ are inferred from our model calculation.

6. DISCUSSION

The kinematics of the F1-4 HII complex and the discrete HII regions in Sgr B2 Main are shown by the radial velocity distribution determined from the first moment images of the H26 α line in Figure 4. The observed kinematics imply certain dynamical processes in the ionized gas associated with the newly born massive stars in the complex. Most of the H26 α line emission arises from the high-density component (A) in an H26 α source in Sgr B2

Main, as shown with our model fitting (see Table 3). The high-density, hyper-compact components with a typical size of a few $\times 10^2 \text{ AU}$ are candidates for rotating ionized disks.

6.1. Candidate for rotating disk

H26 α -7 in F10.39, a weak stimulated source, shows a radial gradient, nearly north-to-south, with radial velocity difference of 9 km s^{-1} or $V_r \approx 4.5 \text{ km s}^{-1}$ across the central $0.2''$, or $2R = 1600 \text{ AU}$, which could indicate an edge-on rotating disk (see Figure 4). If the velocity difference is due to Keplerian rotation of the ionized gas in a disk with an inclination angle i , then the enclosed mass is

$$M \approx 1.1 M_{\odot} \left(\frac{V_r}{\text{km s}^{-1}} \right)^2 \left(\frac{R}{10^3 \text{ AU}} \right) \sin^{-2}(i) \approx 18 \sin^{-2}(i) M_{\odot}, \quad (14)$$

which is consistent with the mass of a late O-type star as the ionizing source. However, we note that the observed kinematics of the two sources can also be interpreted as a rotating-expanding HII shell. The data are not adequate to discriminate between disk and rotating-expanding shell models.

6.2. Confined HII outflow and expanding ionized ring or disk

F3 (H26 α -1) shows a large velocity gradient northwest-to-southeast (see Figure 4). The maximum radial velocity difference is 50 km s^{-1} or $V_r \approx 25 \text{ km s}^{-1}$ across a projected angular size of $0.7''$ or $2R = 5600 \text{ AU}$ which is consistent with the angular size of the ring/disk seen in the VLA high-resolution ($0.049'' \times 0.79''$) observation at 7 mm (De Pree et al. 1998). The ring/disk is nearly circular with a ratio of minor-to-major axis size $\theta_{\text{min}}/\theta_{\text{maj}} \gtrsim 0.5''/0.7''$, indicating an inclination angle $i \lesssim 45^\circ$. If the velocity difference is due to Keplerian rotation of the ionized gas in the ring, then from Equation (14), an enclosed, dynamic mass $1.9 \times 10^3 \sin^{-2}(i) M_{\odot}$ can be inferred, which is at least an order magnitude greater than the inferred total mass of both the ZAMS type O star ($\sim 30 M_{\odot}$) in H26 α -1 and the protostellar core Smm-1 ($\sim 34 M_{\odot}$) in F3 (see Tables 3 and 4). On the other hand, the electron temperature of $T_e = 1.7 \times 10^4 \text{ K}$, inferred for the large, lower-density component (B) from our model fitting, suggests an isothermal sound speed $C_S = \sqrt{kT_e/m_H} = 12 \text{ km s}^{-1}$ which is about a factor of three smaller than the maximum expansion velocity $V_{\text{max}} = V_r \sin^{-1}(i) \approx 35 \text{ km s}^{-1}$. However, the radiation pressure ($P_{\text{rad}} = \frac{L_*}{3c\pi R^2}$) due to a luminosity L_* from an O type star would be comparable to the thermal pressure ($P_{\text{th}} = 2n_e kT_e$) in the HII region with a radius of R , where the factor of 2 accounts for equal densities of electrons (n_e) and protons (n_i),

$$\frac{P_{\text{rad}}}{P_{\text{th}}} = 2.3 \left(\frac{L_*}{10^5 L_{\odot}} \right) \left(\frac{R}{10^3 \text{ AU}} \right)^{-2} \times \left(\frac{n_e}{10^6 \text{ cm}^{-3}} \right)^{-1} \left(\frac{T_e}{10^4 \text{ K}} \right)^{-1}, \quad (15)$$




FIG. 4.— The distribution of intensity weighted radial velocity with respect to a LSR velocity of 58 km s^{-1} is overlaid on the contour image of the continuum emission at 0.86 mm (see Figure 1). The velocity image was constructed from the $\text{H26}\alpha$ line image cube convolved with a circular beam of $0.3''$ using the moment 1 algorithm in Miriad with 4σ intensity cutoff. The contours are $(0.25, 0.35, 0.45, \dots, 0.95) \times 2.5 \text{ Jy beam}^{-1}$ with a circular beam of $0.3''$. The cross signs mark the peak positions of the $\text{H26}\alpha$ line sources of 1, 2, 3, 4 and 6 in the central F1-4 complex (the main figure) along with the sources of $\text{H26}\alpha$ -5, 7, and 8 in the discrete sources with large angular offsets from the center (the insets). Both the vertical and horizontal coordinates are the offsets (arcsec) from the registration center: R.A.(J2000) = $17:47:20.135$ and Decl.(J2000) = $-28:23:04.53$.

where $\frac{P_{\text{rad}}}{P_{\text{th}}}$ is the ratio of the radiation pressure to the thermal pressure. On the basis of the parameters derived for F3 (see Table 3), $\frac{P_{\text{rad}}}{P_{\text{th}}} \approx 0.4$ and 0.7 for the high-density component (A) and the lower-density component (B), respectively. On the other hand, if the radial velocity gradient is mainly due to expansion or outflow, the ram pressure of the expansion motions is $\rho_{\text{HII}} V_{\text{ex}}^2$. Then, the ratio of the ram pressure to the thermal pressure is,

$$\frac{P_{\text{ram}}}{P_{\text{th}}} = 6.1 \times 10^{-3} \left(\frac{T_e}{10^4 \text{ K}} \right)^{-1} \left(\frac{V_{\text{ex}}}{\text{km s}^{-1}} \right)^2. \quad (16)$$

For the lower-density component (B) of F3, $\frac{P_{\text{ram}}}{P_{\text{th}}} \approx 0.5$ if $V_{\text{ex}} \approx C_s$. The derived ratios of $\frac{P_{\text{ram}}}{P_{\text{th}}}$ and $\frac{P_{\text{rad}}}{P_{\text{th}}}$ appear to be consistent. Both the thermal and radiation pressures play a critical role in the expansion of the ionized gas and may accelerate ionized outflow. However both the expansion and the plausible HII outflow are thought to be

confined in the dense ambient medium (De Pree et al. 1998). If the UC HII region is in pressure equilibrium with the ambient medium (Xie et al. 1996), we have,

$$2kn_e T_e \xi = n_{\text{H}} (\mu m_{\text{H}} \sigma_V^2 + kT_k), \quad (17)$$

where σ_V is the turbulent velocity, T_k is the kinetic temperature of the ambient gas, approximately equal to the dust temperature T_d , and $\xi = 1 + \frac{P_{\text{ram}}}{P_{\text{th}}}$ considering the contribution from ram pressure. From Equation (17), we can find a minimum density n_{H} required to confine the HII gas,

$$n_{\text{H}} > 4 \times 10^8 \xi \left(\frac{n_e}{10^6 \text{ cm}^{-3}} \right) \left(\frac{T_e}{10^4 \text{ K}} \right) \left(\frac{\Delta V_{\text{FWHM}}}{\text{km s}^{-1}} \right)^{-2} \text{ cm}^{-3} \quad (18)$$

where $\Delta V_{\text{FWHM}} = \sqrt{(8 \ln 2) \left(\sigma_V^2 + \frac{kT_K}{\mu m_{\text{H}}} \right)}$, the FWHM line width of a molecular line from the surrounding medium. For $T_K \approx T_d = 500 \text{ K}$, the thermal width

is about 3 km s^{-1} . From the SMA measurements of the FWHM line widths for the sixteen kinematic absorption features in $\text{H}_2\text{CO}(3_{30} - 2_{02})$ and $\text{H}_2\text{CO}(3_{31} - 2_{20})$ towards the F1-4 complex (Qin et al. 2008), we find a variance-weighted mean $\Delta V_{\text{FWHM}} = 5 \pm 0.2 \text{ km s}^{-1}$, suggesting a dominant turbulent motion in the line broadening. Thus, from Equation (18) and the parameters derived for the HII regions (Table 3), a minimum $n_{\text{H}} \approx 2 \times 10^7 \text{ cm}^{-3}$ is needed to confine F3 (component B), which is a factor of a few to 10 less than the densities inferred for the protostellar cores but is in good agreement with the value of n_{H} derived from the isothermal model using the observed $T_{\text{b}} = 14 \text{ K}$ of the lowest contour in Figure 4. Our results agree with the predicted range of the molecular n_{H} between $10^7 - 10^8 \text{ cm}^{-3}$ (De Pree et al. 1998).

F1 (H26 α -3, H26 α -4, and H26 α -6) shows complicated kinematics. The line flux from each of the three H26 α sources suggests the presence of at least three O type stars in this region. From analysis of the high-resolution VLA image, De Pree et al. (1998) suggested a total of seven late O-type or early B-type stars in F1. The observed kinematics suggest that complicated dynamics of ionized gas are involved in the OB association.

G (H26 α -8) also shows a broad wing in its H26 α line spectrum, which corresponds to a large radial velocity gradient (ΔV_r) northeast-to-southwest across the source with a size $0.6''$ as seen in the moment 1 image (see Figure 4). Taking the parameters inferred from our modeling (Table 3) and $L_*(\text{O8.5}) \approx 5.4 \times 10^4 L_{\odot}$, with Equation (15), we find that $\frac{P_{\text{rad}}}{P_{\text{th}}} \approx 1$ and ≈ 0.6 for the hypercompact component (A) and the larger component (B), respectively. The observed velocity gradient in the component B region appears to be mainly due to a bipolar ionized outflow accelerated in the region surrounding an O8.5 star. The structure of kinematics for the hypercompact component (A) with a weak stimulated source as suggested earlier was not resolved in these observations.

F10.37 (H26 α -5) shows a broad H26 α line profile with $\Delta V_{\text{FWHM}} = 51 \pm 6 \text{ km s}^{-1}$ but no obvious radial velocity gradient is seen in the moment 1 image (see Figure 4). For $T_e = 1 \times 10^4 \text{ K}$, the thermal broadening gives $\Delta V_{\text{FWHM}}^{\text{th}} = 21 \text{ km s}^{-1}$. Subtracting the thermal broadening, the residual line width $\Delta V_{\text{FWHM}}^{\text{nth}} = 46 \text{ km s}^{-1}$ requires large nonthermal motions. A bipolar ionized outflow along the line-of-sight might be responsible for the large velocity dispersion $\sigma_v = \frac{1}{\sqrt{8 \ln 2}} \Delta V_{\text{FWHM}}^{\text{nth}} = 20 \text{ km s}^{-1}$ in this region.

6.3. Formation of massive protostellar cores

In addition to the H26 α sources, there are about two dozen protostellar cores with masses much greater than the corresponding Bonnor-Ebert masses, and a total molecular mass in the range $5 - 223 M_{\odot}$. The observed filamentary structures in the continuum emission at the submillimeter wavelength suggest that the massive stars and protostellar cores were formed through fragmentation during the initial collapse of the massive core in Sgr B2

Main (Qin et al. 2008). The inferred surface density of each protostellar core appears to be greater than unity; a minimum mass for forming a massive star in order to avoid further fragmentation (Krumholz & McKee 2008). Through competitive accretion, each of the protostellar cores will grow by adding an accreted mass $\dot{M} t_{\text{ff}}$ within the free-fall timescale t_{ff} . The accretion rate can be estimated for an isothermal flow in which the density, $\rho(r) \propto r^{-3/2}$, and the enclosed mass within a radius r , $M(r) \propto r^{3/2}$, can be described by a power law (Shu 1977). The accretion rate is linearly proportional to the accretion velocity V_{Acc} and the enclosed mass M_{R} within $r = R$ but inversely proportional to R ,

$$\dot{M} = V_{\text{Acc}} \frac{dM(r)}{dr} \Big|_{r=R} = 0.3 M_{\odot} \text{ y}^{-1} \left(\frac{V_{\text{Acc}}}{\text{km s}^{-1}} \right) \left(\frac{M_{\text{R}}}{M_{\odot}} \right) \left(\frac{R}{\text{AU}} \right)^{-1}. \quad (19)$$

Based on our measurements $2R \approx \sqrt{5'' \times 3''} \approx 3 \times 10^4 \text{ AU}$, $M_{\text{R}} \sim 1.2 \times 10^3 M_{\odot}$, and $V_{\text{Acc}} \sim C_{\text{S}} = \sqrt{\frac{k T_{\text{K}}}{\mu m_{\text{H}}}} \approx 1.3 \text{ km s}^{-1}$ for a kinetic temperature $T_{\text{K}} \approx T_{\text{d}} = 500 \text{ K}$, we inferred $\dot{M} \approx 0.03 M_{\odot} \text{ y}^{-1}$. For a constant accretion rate of $0.03 M_{\odot} \text{ y}^{-1}$, in a mean free-fall time-scale $t_{\text{ff}} \approx 2 \times 10^3 \text{ y}$, about $60 M_{\odot}$, or $\sim 5\%$ of the total mass increases in the protostellar cores.

We have observed a snapshot of the process of forming a massive-star cluster. A few points regarding the formation of massive star cores can be drawn from our analysis of the SMA observations. First, Figure 4 shows a noticeable angular offset of $\sim 0.1''$ between the H26 α sources (H26 α -1, H26 α -4 and H26 α -6) from their continuum counterparts (Smm-1, Smm-4 and Smm-10). The relative angular offsets are significant ($> 5\sigma$), corresponding to a projected distance of 800 AU. If the H26 α sources denote the locations of the new born O stars, the offsets of the protostellar cores (Smm-1, Smm-4 and Smm-10) from the ionizing stars might suggest that the local accretion centers are altered after the onset of the radiation from the O stars. A massive protostellar core in the immediate vicinity of a UC HII region has been also observed in the W51e2 complex (Shi, Zhao, & Han 2010a,b). The authors use SMA high-resolution observations to show that the protostellar core (W51e2-E) with a mass of $140 M_{\odot}$, which is located $0.9''$ (4600 AU) from the UC HII region (W51e2-W), becomes the new accretion center and undergoes active star formation ejecting a powerful molecular outflow. Multiple pairs of O-type star and massive protostellar core found in Sgr B2 Main show a cluster version of the W51e2 system. Consequently, a considerable amount of mass must have existing in each of the paired protostellar cores before the O-type stars were born unless there were large accretion rates, $\sim 0.1 M_{\odot} \text{ s}^{-1}$ at each of the massive protostellar cores in the past 1×10^3 years after the onset of radiation from the O stars. The protostellar cores and their companion massive stars likely originate from fragmentation during the initial collapse of the cloud. After the primitive molecular clumps formed, the massive ones undergo substantial gravitational collapse to form stars. Competitive accretion may occur in the process of forming a cluster of massive stars in a massive core like Sgr B2 Main. How-

ever, from our observations of this particular phase of massive star formation in a cluster, it is uncertain how much mass has been added to each of the protostellar cores through competitive accretion owing to the uncertainty in the time-scale of the process. On the basis of our analysis, accretion will add a small fraction of mass to each of the protostellar cores in a mean free-fall time-scale of 2×10^3 y.

7. CONCLUSION AND SUMMARY

Using SMA archival data observed in 2007 for Sgr B2 Main, we imaged the H26 α line and continuum emission at 0.86 mm with natural and uniform weighted synthesized beams of $0.40'' \times 0.28''$ and $0.36'' \times 0.22''$, respectively. Eight compact H26 α line sources have been detected towards the central $5''$ (~ 0.2 pc). Using the non-LTE approach, we fitted the H26 α line along with other available RRL and radio continuum data with two isothermal, homogeneous HII components: 1) high-density (a few $\times 10^7$ cm $^{-3}$), hyper-compact (several hundred AU) and 2) lower-density (a few $\times 10^6$ cm $^{-3}$), ultra-compact (a few thousand AU). The observed H26 α line fluxes suggest that each of the eight H26 α line emission regions must be powered by a Lyman continuum source from an O type star or a cluster of B-type stars. The typical age of the HII regions are a thousand years. The brightest H26 α -1 source in F3 appears to be associated

with an O6 star. The H26 α line from H26 α -7 (F10.39) appears to be enhanced by stimulated emission in the vicinity of an O9.5 star.

About two dozen compact continuum emission cores are detected at 0.86 mm in the central $5''$ (0.2 pc). Using a power-law distribution of dust temperature and observed brightness temperature, we determined the physical properties for each of the protostellar cores with masses in the range of 5 to $\gtrsim 200$ solar masses. The surface densities of the protostellar cores are in the range 13 to 150 g cm $^{-2}$, and molecular densities 1×10^8 to 1×10^9 cm $^{-3}$. We also calculated the Bonnor-Ebert mass (M_{BE}) for each core. The total molecular mass of each core appears to be much greater than the value of M_{BE} , suggesting that the cores undergo a substantial gravitational collapse to form massive stars with a typical free-fall time scale of a few thousand years.

We are grateful to the anonymous referee for thoughtful comments and suggestions which were helpful for improving this paper. We thank Dr. Hui Shi for her assistance preparing Miriad scripts for the reduction of the SMA data. JHZ is grateful to the National Astronomical Observatories of China for hosting his visit during the course of writing this research paper.

APPENDIX

LYMAN CONTINUUM PHOTON FLUX

In this Appendix, we show the detailed procedure to derive the Lyman continuum photon flux from the measurements of a hydrogen radio recombination line.

Photoionizations and recombinations of hydrogen in HII regions

For an HII region with electron density n_e , the ionization equilibrium can be described by the balance between photoionizations and recombinations of electrons with the ions. The number of ionizing photons (N_I) emitted by a star in unit time, or the Lyman continuum photon flux (N_{Lym}), is equal to the number of recombinations ($N_R = \frac{4\pi}{3} R_s^3 n_e^2 \alpha_B$) to the excited levels within the Strömgren radius (R_s) in unit time (Osterbrock 1974):

$$N_{Lym} = \frac{4\pi}{3} R_s^3 n_e^2 \alpha_B \quad (A1)$$

where $\alpha_B = \alpha_B(T_e)$ is the recombination coefficient in case B, which is a function of electron temperature (T_e) with a value $\alpha_B = 2.58 \times 10^{-13}$ cm $^{-3}$ s $^{-1}$ at $T_e = 1 \times 10^4$ K (Hummer & Seaton 1963); and $R_s^3 \approx \frac{3}{4\pi} V \approx \frac{3}{4\pi} \Omega_{HII} D^2 L f_V$.

For an HII region located at distance D , with solid angle Ω_{HII} , path length L , and electron volume filling factor f_V , Equation (A1) can be rewritten as,

$$N_{Lym} = n_e^2 L f_V \alpha_B \Omega_{HII} D^2 \quad (A2)$$

Optical depth and flux density of hydrogen radio recombination lines

For an isothermal-homogeneous HII region, the optical depths for hydrogen radio recombination lines (τ_L) and free-free continuum (τ_C), are proportional to the emission measure ($EM = n_e^2 L f_V$), and are given by Shaver (1975):

$$\tau_L = b_N \beta_N \tau_L^* \approx 575 b_N \beta_N \left(\frac{\nu}{\text{GHz}} \right)^{-1} \left(\frac{n_e}{\text{cm}^{-3}} \right)^2 \left(\frac{L f_V}{\text{pc}} \right) \left(\frac{T_e}{\text{K}} \right)^{-5/2} \left(\frac{\Delta V_D}{\text{km s}^{-1}} \right)^{-1} \left(1 + 1.48 \frac{\Delta V_P}{\Delta V_D} \right)^{-1}, \quad (A3)$$

and

$$\tau_C \approx 0.08235 \left(\frac{n_e}{\text{cm}^{-3}} \right)^2 \left(\frac{L f_V}{\text{pc}} \right) \left(\frac{\nu}{\text{GHz}} \right)^{-2.1} \left(\frac{T_e}{\text{K}} \right)^{-1.35} a(\nu, T_e), \quad (A4)$$

where τ_L^* is the LTE line optical depth; b_N and β_N are the population departure coefficients; n_e is the electron density; ΔV_D and ΔV_P are the FWHM Doppler and pressure line widths in km s^{-1} , respectively. The correction factor $a(\nu, T_e)$ is of order unity.

On the other hand, if an HII region, with no dust in it, is optically thin and LTE ($\tau_C \ll 1$, $\tau_L^* \ll 1$, and $b_N = \beta_N = 1$), the line flux density from Equation (1) in the main text becomes,

$$S_L^* \approx \frac{2k\nu^2}{c^2} \Omega_{\text{HII}} T_e \tau_L^* \quad (\text{A5})$$

Normally, the Lyman continuum photon flux (N_{Lyman}) is derived from the emission measure determined from the free-free continuum flux density observed at a radio frequency, *e.g.* Wilcots (1994). In the following, we show the derivation of the emission measure from a radio recombination line flux. Therefore, N_{Lyman} can be determined directly from a radio recombination line flux.

Emission measure from hydrogen radio recombination lines

From Equations (A3) and (A5), neglecting the pressure broadening ($\Delta V_D \gg \Delta V_P$) the emission measure for an optically-thin, LTE gas can be expressed as,

$$n_e^2 L f_V = 5.66 \times 10^{-8} \text{cm}^{-6} \text{pc} \left(\frac{S_L^* \Delta V_D}{\text{Jy km s}^{-1}} \right) \left(\frac{\nu}{\text{GHz}} \right)^{-1} \left(\frac{T_e}{\text{K}} \right)^{3/2} \Omega_{\text{HII}}^{-1} \quad (\text{A6})$$

For a general isothermal, homogeneous HII region, the emission measure can be determined by the following equation,

$$n_e^2 L f_V = 5.66 \times 10^{-8} \text{cm}^{-6} \text{pc} \left(\frac{S_L \Delta V_D}{\text{Jy km s}^{-1}} \right) \left(\frac{\nu}{\text{GHz}} \right)^{-1} \left(\frac{T_e}{\text{K}} \right)^{3/2} \Omega_{\text{HII}}^{-1} \Psi_N^{-1} \quad (\text{A7})$$

where Ψ_N is the ratio of the non-LTE line flux (S_L) to the LTE line flux in optically thin ($S_L^*[\tau_L^* \ll 1]$) for the line transition $N+1 \rightarrow N$, where N is the principal quantum number. Ψ_N is a function of the line and continuum optical depths (τ_L and τ_C) and departure coefficients (b_N and β_N),

$$\Psi_N \equiv \frac{S_L}{S_L^*[\tau_L^* \ll 1]} \approx \left[\left(\frac{\tau_L/\beta_N + \tau_C}{\tau_L + \tau_C} \right) (1 - e^{-(\tau_L + \tau_C)}) - (1 - e^{-\tau_C}) \right] \tau_L^{*-1}, \quad (\text{A8})$$

neglecting the term associated with the dust emission (S_d) in Equation (1) of the main text.

Lyman continuum photon flux from hydrogen radio recombination lines

For an electron temperature (T_e) in an isothermal, homogeneous HII region, inserting $n_e^2 L f_V$ of Equation (A7) into Equation (A2), we derived the Lyman continuum photon flux from the measurements with flux density (S_L) and FWHM of Doppler broadening (ΔV_D) for a hydrogen radio recombination line ($N+1 \rightarrow N$) at frequency ν_L ,

$$N_{\text{Lyman}} = 4.29 \times 10^{47} \left(\frac{S_L \Delta V_D}{\text{Jy km s}^{-1}} \right) \left(\frac{\nu_L}{\text{GHz}} \right)^{-1} \left(\frac{D}{\text{kpc}} \right)^2 \left(\frac{T_e}{10^4 \text{K}} \right)^{1.5} \left(\frac{\alpha_B(T_e)}{\alpha_B(10^4 \text{K})} \right) \Psi_N^{-1} \text{ ph s}^{-1} \quad (\text{A9})$$

where Ψ_N corrects for the effects due to the optical depths and non-LTE distribution. Thus, from Equation (A8), $\Psi_N \approx 1$ for optically thin, LTE gas; $\Psi_N \approx b_N(1 - \frac{1}{2}\tau_C\beta_N)$ for optically thin, non-LTE gas; and $\Psi_N \approx \frac{e^{-\tau_C}(1 - e^{-\tau_L^*})}{\tau_L^*}$ for the LTE gas becomes optically thick.

REFERENCES

- | | |
|---|--|
| <p>Bonnell, I. A., Bate, M. R. & Zinnecker, H., 1998, MNRAS, 298, 93</p> <p>Bonnell, I. A., & Bate, M. R., 2002, MNRAS, 336, 659</p> <p>Bonnell, I. A., Vine, S. G. & Bate, M. R., 2004, MNRAS, 349, 735</p> <p>Bonnell, I. A., & Bate, M. R., 2006, MNRAS, 370, 488</p> <p>Bonnell, I. A. 2008, in ASP Conf. Series, Vol. 390: Pathways through an Eclectic Universe, 26</p> <p>Bonnor, W. B. 1956, MNRAS, 116, 351</p> <p>Brocklehurst, M., & Leeman, S. 1971, Astroph. Lett, 9, 35</p> <p>Brocklehurst, M., & Seaton, M. J. 1972, MNRAS, 157, 179</p> <p>Cernicharo, J., Goicoechea, J. R. & Pardo J. R. 2006, ApJ, 642, 940</p> | <p>Churchwell, E. 2002, ARA&A, 40, 27</p> <p>De Pree, C. G., Gaume, R. A., Goss, W. M., & Claussen, M. J. 1996, ApJ, 464, 788</p> <p>De Pree, R. A., Goss, W. M., & C. G., Gaume 1998, ApJ, 464, 788</p> <p>Ebert, R. 1957, Z. Astrophys., 42, 263</p> <p>Figer, D. F., 2005, Nature, 434, 192</p> <p>Figer, D. F., 2008, in IAU proc. 250, Massive Stars as Cosmic Engines, eds. Bresolin, Crowther & Puls, 247</p> <p>Garay, G. & Lizano, S. 1999, PASP, 111, 1049</p> <p>Gaume, R. A. & Claussen, M. J. 1990, ApJ, 351, 538</p> |
|---|--|

- Gaume, R. A., Claussen, M. J., De Pree, C. G., Goss, W. M., & Mehringer, D. W. 1995, *ApJ*, 449, 663
- Goldsmith, P. F. & Lis, D. C. 1990, *ApJ*, 350, 186
- Goldsmith, P. F. & Lis, D. C., Lester, D. F., & Harvey, P. M., 1992, *ApJ*, 389, 338
- Gordon, M. A., & Sorooshenko, R. L. 2002, *Radio Recombination Lines. Their Physics and Astronomical Applications* (Dordrecht: Kluwer)
- Hummer, D. G. & Seaton, M. J. 1963, *MNRAS*, 125, 437
- Kahn, F. D., 1974, *A&A*, 37, 149
- Krumholz M. R., McKee, C. F. & Klein, R. I., 2005, *Nature*, 438, 332
- Krumholz M. R. & McKee, C. F. 2008, *Nature*, 451, 28
- Krumholz, M. R., Klein, R. I., McKee, C. F., Offner, S. S. R. & Cunningham, A. J., 2009, *Science*, 323, 754
- Krumholz, M. R., Cunningham, A. J., Klein, R. I., & McKee, C. F. 2010, *ApJ*, 713, 1120
- Larson, R. B. & Starrfield, S., 1971, *A&A*, 13, 190
- Lis, D. C. & Goldsmith, P. F. 1990, *ApJ*, 356, 195
- Liu, S.-Y. & Snyder, L. E. 1999, *ApJ*, 523, 683
- Mathis, J. S., Rumpl, W. & Nordsieck, K. H., 1977, *ApJ*, 217, 425
- McKee, C. F. & Tan, J. C., 2002, *Nature*, 416, 59
- McKee, C. F. & Ostriker, E. C., *ARA&A*, 45, 565
- McLaughlin, D. E. & Pudritz, R. E., 1997, *ApJ*, 476, 750
- Nakano, T., 1989, *ApJ*, 345, 464
- Osoiro, M., Lizano, S. & D'Alessio, P. 1999, *ApJ*, 525, 808
- Ossenkopf, & Henning, T. 1994, *A&A*, 291, 943
- Osterbrock, D. E. 1974, *Astrophysics of Gaseous Nebulae*, (San Francisco: W. H. Freeman and Company)
- Panagia, N. 1973, *AJ*, 78, 929
- Qin, S.-L., Zhao, J.-H., Moran, J. M., Marrone, D. P., Patel, N. A., Wang, J.-J., Liu, S.-Y. & Kuan, Y.-J., 2008, *ApJ*, 677, 353
- Reid, M. J., Menten, K. M., Zheng, X. W., Brunthaler, A., & Xu, Y., 2009, *ApJ*, 705, 1548
- Rolfs, R. et al. 2010, *A*, 521, L46
- Sault, R. J., Teuben, P. J., & Wright, M. C. H. 1995, in *Astronomical Society of the Pacific Conference Series*, Vol. 77, *Astronomical Data Analysis Software and Systems IV*, ed. R. A. Shaw, H. E. Payne, & J. J. E. Hayes, 433–+
- Scoville, N. Z. & Kwan, J. 1976, *ApJ*, 206, 718
- Shaver, P. A. 1975, *Pramana*, 5, 1
- Shi, H., Zhao, J., & Han, J. L. 2010a, *ApJ*, 710, 843
- Shi, H., Zhao, J., & Han, J. L. 2010b, *ApJ*, 718, L181
- Shu, F. 1977, *ApJ*, 214, 488
- Spitzer, L., 1978, *Physical Processes in the Interstellar Medium* (New York: Wiley)
- Strömgren, B. 1939, *ApJ*, 89, 526
- Vacca, W. D., Garmany, C. D., & Shull, J. M. 1996, *ApJ*, 460, 914
- Walmsley, C. M. 1990, *A&A Suppl. Ser.*, 82, 201
- Wilcots, E. M., 1994 *AJ*, 107, 1338
- Wolfire, M. G. & Cassinelli, J. P., 1986, *ApJ*, 310, 207
- Wolfire, M. G. & Cassinelli, J. P., 1987, *ApJ*, 319, 850
- Xie, T., Mundy, L. G., Vogel, S. N., & Hofner, P. 1996, *ApJ*, 473, L131
- Zhao, J.-H., Blundell, R., Moran, J. M., Downes, D., Fchuster, K. F., & Marrone, D. 2010, *ApJ*, 723, 1097

TABLE 1
LOG OF OBSERVATIONS AND IMAGING

Parameters	H26 α	Continuum
OBSERVATION:		
Date	2007 June 18	
Array configuration	Very extended (8 antennas)	
Pointing R.A. (J2000)	17:47:20.151	
Pointing decl. (J2000)	-28:23:04.79	
ν_{LO}	348.664 GHz	
$\delta\nu$	0.406 MHz	
Bandwidth	2GHz(USB) & 2GHz(LSB)	
On-source time	1.95 hr	
T_{sys}	150-500 K	
CALIBRATORS:		
Flux density	Callisto	
Phase	J1733-130	
Bandpass	J1229+020, J1733-130, J1743-038, J1751+096, J2015+371	
IMAGING:		
ν_0 or λ	353.623 GHz	0.86 mm
channel width	3 km s ⁻¹	...
FWHM beam	0.40'' $\times$ 0.28''(15°)	0.36'' $\times$ 0.22''(13°)
r.m.s.	0.1 Jy bm ⁻¹ chan ⁻¹	8 mJy bm ⁻¹

TABLE 2
SUBMILLIMETER PROPERTIES OF SGR B2 (MAIN)

Region	The H26 α Line Measurements						
	$\Delta\alpha^a$ (")	$\Delta\delta^a$ (")	$\theta_{Maj} \times \theta_{Min}, PA$ (" $\times$ " , $^\circ$)	$S_{H26\alpha}$ (Jy)	V_{LSR} (km s $^{-1}$)	ΔV_{FWHM} (km s $^{-1}$)	S_C (Jy)
H26 α -1 (F3)	-0.17	+0.03	0.29 $\times$ 0.21, -25	7.0 $\pm$ 0.1	64 $\pm$ 1	40 $\pm$ 1	2.9 $\pm$ 0.5
H26 α -2 (F4)	+0.37	+0.37	U	0.63 $\pm$ 0.08	66 $\pm$ 2	35 $\pm$ 8	0.41 $\pm$ 0.10
H26 α -3 (F1A)	-0.53	+0.73	U	1.2 $\pm$ 0.1	48 $\pm$ 2	54 $\pm$ 5	1.8 $\pm$ 0.4
H26 α -4 (F1B)	-0.92	+0.38	U	0.8 $\pm$ 0.1	73 $\pm$ 1	36 $\pm$ 4	1.0 $\pm$ 0.4
H26 α -5 (F10.37)	-0.00	-1.18	U	0.62 $\pm$ 0.08	52 $\pm$ 2	51 $\pm$ 6	0.77 $\pm$ 0.10
H26 α -6 (F10.303)	-0.90	+1.08	U	0.74 $\pm$ 0.1	57 $\pm$ 2	50 $\pm$ 6	0.91 $\pm$ 0.24
H26 α -7 (F10.39)	+0.22	-1.97	U	0.62 $\pm$ 0.09	69 $\pm$ 2	29 $\pm$ 6	0.35 $\pm$ 0.09
H26 α -8 (G)	+1.45	+1.70	U	0.73 $\pm$ 0.1	53 $\pm$ 2	48 $\pm$ 6	0.2 $\pm$ 0.1

Source	The Continuum Measurements at 0.8 mm						
	$\Delta\alpha^a$ (")	$\Delta\delta^a$ (")	$\theta_{Maj} \times \theta_{Min}, PA$ (" $\times$ " , $^\circ$)	S_P (Jy beam $^{-1}$)	S_T (Jy)	$\Delta\theta^a$ (")	T_b (10 ^{2} K)
Smm-1	-0.20	+0.16	0.80 $\times$ 0.40, +10	2.6 $\pm$ 0.2	11 $\pm$ 0.8	0.26	3.4
Smm-2	+0.78	+0.09	0.22	0.4 $\pm$ 0.1	0.4 $\pm$ 0.1	0.79	0.8
Smm-3	-0.39	-0.72	0.35 $\times$ 0.25, +3	2.2 $\pm$ 0.2	2.2 $\pm$ 0.2	0.82	2.5
Smm-4	-0.49	+0.68	0.83 $\times$ 0.30, +32	2.0 $\pm$ 0.3	7.0 $\pm$ 0.9	0.84	2.7
Smm-5	+1.00	+0.41	0.22	0.4 $\pm$ 0.1	0.4 $\pm$ 0.1	1.08	0.8
Smm-6	-0.92	+0.59	0.35 $\times$ 0.26, +21	1.1 $\pm$ 0.2	2.2 $\pm$ 0.3	1.09	2.4
Smm-7	+0.01	-1.20	0.61 $\times$ 0.31, -31	0.9 $\pm$ 0.1	2.7 $\pm$ 0.3	1.20	1.4
Smm-8	-0.68	-1.09	0.22	0.8 $\pm$ 0.2	0.8 $\pm$ 0.2	1.28	1.6
Smm-9	+0.58	-1.17	0.30 $\times$ 0.15, -75	0.5 $\pm$ 0.1	0.6 $\pm$ 0.1	1.31	2.0
Smm-10	-1.00	+1.01	0.38 $\times$ 0.22, -21	1.3 $\pm$ 0.2	1.7 $\pm$ 0.2	1.42	2.0
Smm-11	+1.03	+1.04	0.22	0.3 $\pm$ 0.1	0.3 $\pm$ 0.1	1.46	0.6
Smm-12	-0.08	+1.59	0.53 $\times$ 0.14, -82	0.4 $\pm$ 0.1	0.8 $\pm$ 0.2	1.59	1.1
Smm-13	-0.77	-1.57	0.53 $\times$ 0.45, -75	1.2 $\pm$ 0.2	3.9 $\pm$ 0.6	1.75	1.6
Smm-14	+0.33	-1.74	0.39 $\times$ 0.19, -48	0.8 $\pm$ 0.2	1.3 $\pm$ 0.3	1.77	1.7
Smm-15	-1.77	-0.02	0.45 $\times$ 0.09, +29	0.4 $\pm$ 0.2	0.4 $\pm$ 0.2	1.77	1.0
Smm-16	+0.83	-1.67	0.22	0.4 $\pm$ 0.1	0.4 $\pm$ 0.1	1.86	0.8
Smm-17	-0.89	+1.64	0.53 $\times$ 0.20, -13	0.8 $\pm$ 0.1	1.6 $\pm$ 0.2	1.87	1.5
Smm-18	-1.77	+1.24	0.22	0.4 $\pm$ 0.1	0.4 $\pm$ 0.1	2.16	0.8
Smm-19	+0.53	-2.12	0.22	0.7 $\pm$ 0.1	0.7 $\pm$ 0.1	2.19	1.4
Smm-20	-2.12	+0.69	0.22	0.4 $\pm$ 0.1	0.4 $\pm$ 0.1	2.23	0.8
Smm-21	+1.46	+1.74	0.22	0.2 $\pm$ 0.1	0.2 $\pm$ 0.1	2.27	0.4
Smm-22	-1.57	+2.04	0.30 $\times$ 0.19, +27	0.5 $\pm$ 0.2	0.5 $\pm$ 0.2	2.57	0.9
Smm-23	+0.56	-2.57	0.60 $\times$ 0.22, +57	0.4 $\pm$ 0.1	1.0 $\pm$ 0.1	2.63	0.7

^a Offsets from F3 at RA(J2000)=17:47:20.178, Decl(J2000)=-28:23:04.67

Table 3. Hyper- and Ultra-Compact HII cores in Sgr B2 (Main)

	H26 α -1 (F3)		H26 α -2 (F4)		H26 α -3 (F1-A)		H26 α -4 (F1-B)		H26 α -5 (F10.37)		H26 α -6 (F10.303)		H26 α -7 (F10.39)		H26 α -8 (G)	
	A	B	A	B	A	B	A	B	A	B	A	B	A	B	A	B
R (10^3 AU)	2.0	4.0	0.8	3.2	1.0	4.0	0.4	2.7	0.61	2.4	0.91	2.8	0.16	1.2	0.4	2.8
HII GAS PROPERTIES																
T_e (K)	9.0	17	10	10	9.5	14	5.0	10	5.0	10	5.0	10	7.0	10	6.5	10
n_e (10^5 cm $^{-3}$)	42	3.2	50	2.0	52	3.1	85	0.1	52	1.5	30	2.5	330	3.5	110	2.6
f_V	0.1	0.5	0.1	0.5	0.1	0.5	0.1	0.5	0.1	0.5	0.1	0.5	0.1	0.5	0.1	0.5
b_{26}	0.91	0.82	0.92	0.73	0.92	0.79	0.93	0.67	0.90	0.72	0.86	0.74	0.97	0.76	0.94	0.74
β_{26}	-7.4	-13.8	-7.0	-11.2	-6.7	-13.6	-3.1	-4.2	-4.5	-10.0	-6.3	-12.0	-1.2	-13.1	-3.1	-12.2
τ_L (H26 α)	-0.60	-0.008	-0.30	-0.012	-0.33	-0.015	-1.1	$\sim$ 0.0	-0.61	-0.003	-0.36	-0.011	-1.3	-0.02	-0.71	-0.014
τ_C (0.86mm)	0.024	<0.001	0.012	<0.001	0.019	0.001	0.040	$\sim$ 0.0	0.023	<0.001	0.011	<0.001	0.16	<0.001	0.050	<0.001
$\zeta_{H26\alpha}$	94	6	73	27	71	29	99	1	94	6	81	19	95	5	81	19
Ψ_{26}	1.3	0.82	1.1	0.73	1.1	0.80	1.7	0.67	1.3	0.72	1.1	0.74	2.0	0.77	1.4	0.75
τ_L (H66 α)	0.14	-0.039	0.058	-0.085	0.093	-0.07	0.71	-0.001	0.036	-0.033	0.019	-0.068	0.37	-0.087	0.32	-0.073
τ_C (13mm)	9.5	0.25	4.8	0.15	7.3	0.30	4.2	$\sim$ 0.0	9.8	0.065	1.2	0.052	67	0.044	20	0.23
EM (10^{10} cm $^{-6}$ pc)	1.6	0.10	0.97	0.03	1.4	0.093	1.4	<0.001	0.80	0.013	0.40	0.042	8.4	0.035	2.3	0.045
M_{HII} ($10^{-2} M_\odot$)	1.0	3.3	0.081	1.0	0.19	3.2	0.017	0.030	0.038	0.33	0.072	0.85	<0.005	0.1	0.022	0.88
U (pc cm $^{-2}$)	74	47	34	27	45	46	24	3	27	17	28	27	24	15	29	29
$\log[N_{Ly\alpha}(ph\ s^{-1})]$	49.16	48.31	48.10	47.81	48.50	48.34	47.90	44.98	48.03	47.19	48.07	47.82	47.77	47.02	48.04	47.85
$\log[N_{Ly\alpha}^{A+B}(ph\ s^{-1})]$	49.22		48.28		48.73		47.90		48.09		48.26		47.84		48.26	
ZAMS-type	O6		O8.5		O7		O9.5		O9		O8.5		O9.5		O8.5	
t_{age} (y)	1.3		1.1		1.1		0.9		0.8		0.9		0.4		0.9	
DUST PROPERTIES ($T_d = 500$ K, $\beta = 1.5$)																
τ_d (0.86mm)	0.11		0.03		0.11		0.17		0.15		0.12		0.32		<0.01	
M_{tot} ($M_\odot$)	11		1.9		11		7.5		5.4		5.9		2.9		<0.01	

TABLE 4
PROPERTIES OF PROTOSTELLAR CORES IN SGR B2 (MAIN)

Name	r (pc)	size (10^3 AU)	T_d (10^2 K)	τ_d	M ($M_\odot$)	Σ ($g\text{ cm}^{-2}$)	$n(\text{H})$ (10^8 cm^{-3})	M_{BE} ($M_\odot$)	t_{ff} (10^3 yr)
Smm-1	0.01	4.5	5.0	1.1	143	55	2.1	2.9	2.3
Smm-2	0.05	1.8	2.5	0.4	8	20	1.9	1.0	2.4
Smm-3	0.04	2.4	2.6	2.6	93	130	9.5	0.5	1.1
Smm-4	0.04	4.4	2.7	1.9	223	92	3.7	0.9	1.8
Smm-5	0.07	1.8	2.2	0.4	9	22	2.2	0.9	2.3
Smm-6	0.05	2.4	2.5	3.0	111	150	10.8	0.4	1.0
Smm-7	0.07	3.5	2.2	1.0	74	48	2.4	0.8	2.2
Smm-8	0.07	1.8	2.2	1.3	25	63	6.2	0.5	1.4
Smm-9	0.08	1.7	2.1	0.9	17	46	4.8	0.5	1.6
Smm-10	0.07	2.3	2.2	2.3	77	112	8.4	0.4	1.2
Smm-11	0.09	1.8	2.1	0.3	7	17	1.7	0.9	2.6
Smm-12	0.09	2.2	2.1	0.7	21	35	2.8	0.7	2.0
Smm-13	0.09	3.9	2.0	1.6	155	79	3.5	0.6	1.8
Smm-14	0.10	2.2	1.9	2.1	64	105	8.4	0.4	1.2
Smm-15	0.09	1.6	2.0	0.6	10	32	3.4	0.6	1.8
Smm-16	0.11	1.8	1.9	0.6	11	27	2.7	0.6	2.1
Smm-17	0.10	2.6	2.0	1.4	59	68	4.5	0.5	1.6
Smm-18	0.11	1.8	1.9	0.6	11	28	2.7	0.6	2.1
Smm-19	0.12	1.8	1.8	1.5	30	75	7.4	0.3	1.2
Smm-20	0.11	1.8	1.9	0.6	11	28	2.8	0.6	2.0
Smm-21	0.13	1.8	1.8	0.3	5	13	1.3	0.8	3.0
Smm-22	0.13	1.9	1.8	0.7	15	33	3.0	0.5	2.0
Smm-23	0.15	2.9	1.7	0.6	31	28	1.7	0.6	2.6